

BOTOX[®]
onabotulinumtoxinA injection
CHRONIC MIGRAINE

“What can I expect with BOTOX[®]?”

An effective preventive treatment in Chronic Migraine.

BOTOX[®] prevents on average 8 to 9 headache days and migraine/probable migraine days a month (vs 6 to 7 for placebo)



Congrats on taking this exciting next step! This folder is your guide to starting and getting the most from BOTOX[®] for Chronic Migraine.

BOTOX[®]
SAVINGS
PROGRAM

You may pay

\$0 for treatment
See how on page 6.

By participating in the BOTOX[®] Savings Program, you acknowledge and agree to the full Terms & Conditions set out at BOTOXSavingsProgram.com/TermsandConditions. Patients enrolled in Medicare, Medicaid, TRICARE, or any other government reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.



Indication

BOTOX[®] is a prescription medicine that is injected to prevent headaches in adults with chronic migraine who have 15 or more days each month with headache lasting 4 or more hours each day in people 18 years or older.

It is not known whether BOTOX[®] is safe and effective to prevent headaches in patients with migraine who have 14 or fewer headache days each month (episodic migraine).

IMPORTANT SAFETY INFORMATION

BOTOX[®] may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX[®]:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months

- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, loss of bladder control, trouble breathing, and trouble swallowing

There has not been a confirmed serious case of spread of toxin effect away from the injection site when BOTOX[®] has been used at the recommended dose to treat chronic migraine.

BOTOX[®] may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of taking BOTOX[®]. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Please see additional Important Safety Information inside.



“What can I expect with BOTOX®?”


a treatment with a 10+ year LEGACY in Chronic Migraine

BOTOX® has been approved in Chronic Migraine for 10+ years



FIRST APPROVED

In 2010, BOTOX® became the **first preventive treatment** approved by the FDA for Chronic Migraine



MILLIONS OF TREATMENTS

4.2 million BOTOX® treatments have been given to over **600,000 people** living with Chronic Migraine since 2010



PATIENTS STICK WITH IT

In a survey, **97% of Chronic Migraine patients** being treated with BOTOX® say they plan to continue treatment*



Enroll today at [BOTOXSavingsProgram.com](https://www.BOTOXSavingsProgram.com)*

*By participating in the BOTOX® Savings Program, you acknowledge and agree to the full Terms & Conditions set out at [BOTOXSavingsProgram.com/TermsandConditions](https://www.BOTOXSavingsProgram.com/TermsandConditions). Patients enrolled in Medicare, Medicaid, TRICARE, or any other government-reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.

YouTube



Want to hear from real BOTOX® patients? Scan this QR code to see their experience.



*Based on interviews with 71 people currently taking BOTOX®

IMPORTANT SAFETY INFORMATION (continued)

Do not receive BOTOX® if you: are allergic to any of the ingredients in BOTOX® (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as *Myobloc*® (rimabotulinumtoxinB), *Dysport*® (abobotulinumtoxinA), or *Xeomin*® (incobotulinumtoxinA); have a skin infection at the planned injection site.

The dose of BOTOX® is not the same as, or comparable to, another botulinum toxin product.

Please see additional Important Safety Information throughout this brochure.

“What can I expect with BOTOX[®]?”

a treatment that's TESTED and PROVEN



“What can I expect with BOTOX[®]?”



a treatment that's GIVEN with a PURPOSE

PROVEN EXPERIENCE:

A decade of clinical studies paved the way for BOTOX[®] receiving FDA approval in 2010. Since then, BOTOX[®] has built over 10 years of experience helping patients with Chronic Migraine.

PREVENTS HEADACHES/MIGRAINES BEFORE THEY START:

Studies found that BOTOX[®] prevents on average 8 to 9 headache days and migraine/probable migraine days a month (vs 6 to 7 for placebo).

HAS A KNOWN SAFETY PROFILE:


BOTOX[®] studies measured side effects and found the benefits outweigh the risks. The most common side effects were neck pain (9% for BOTOX[®] vs 3% for placebo) and headache (5% vs 3%).*



About 10 minutes of treatment every 3 months is what you can expect



Your doctor will inject BOTOX[®] for Chronic Migraine across 7 head/neck muscle areas that may be linked to migraine (31 injections total)



The **needle** used is a half-inch long and **very thin**. Injections may feel like tiny pinches or pinpricks

BOTOX[®] Treatment Journey

WEEK 0
FIRST TREATMENT
Make your next appointment before you leave the office to stay on schedule

WEEK 4
INITIAL RESULTS
You may start to feel results by week 4, but full effectiveness won't be measured until 24 weeks

WEEK 12
SECOND TREATMENT
Your second treatment should be 12 weeks after the first to get full benefit of treatment

WEEK 24
EVALUATING RESULTS
You and your doctor will see how well BOTOX[®] is working for you after 24 weeks and decide whether to continue treatment every 12 weeks

*Other common side effects include migraine (4% vs 3%), slight or partial facial paralysis (2% vs 0%), drooping eyelids (4% vs <1%), bronchitis (3% vs 2%), musculoskeletal stiffness (4% vs 1%), muscular weakness (4% vs <1%), muscle pain (3% vs 1%), musculoskeletal pain (3% vs 1%), muscle spasms (2% vs 1%), injection-site pain (3% vs 2%), and high blood pressure (2% vs 1%).

IMPORTANT SAFETY INFORMATION (continued)
Serious and/or immediate allergic reactions have been reported including itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX[®] should be discontinued.

Tell your doctor about all your muscle or nerve conditions such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects including difficulty swallowing and difficulty breathing from typical doses of BOTOX[®]

Please see additional Important Safety Information throughout this brochure.



“What can I expect with BOTOX[®]?”

a treatment you can ACCESS



You may pay
\$0 for treatment

By participating in the BOTOX[®] Savings Program, you acknowledge and agree to the full Terms & Conditions set out at BOTOXSavingsProgram.com/TermsandConditions. Patients enrolled in Medicare, Medicaid, TRICARE, or any other government-reimbursed healthcare program are not eligible. Other restrictions and maximum limits apply.

- **≈ 100%** of commercial insurance plans cover BOTOX[®], and most cover the majority of BOTOX[®] costs
- If you're eligible, the BOTOX[®] Savings Program **can help you pay** for BOTOX[®] and its treatment fees by saving you up to \$1000 per treatment*
- **Text SAVE to 27747,**[†] visit BOTOXSavingsProgram.com, or call 1-800-44-BOTOX to get started

*For residents of Massachusetts and Rhode Island, offer applies only to the cost of BOTOX[®] and not to any related medical service(s).

[†]See Privacy & Terms: <http://bit.ly/2RvxiWr>. Message & data rates may apply. Message frequency may vary. Text HELP for help or STOP to end.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all your medical conditions, including if you: have or have had bleeding problems; have plans to have surgery; had surgery on your face; weakness of forehead muscles; trouble raising your eyebrows; drooping eyelids; any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX[®] can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX[®] passes into breast milk).

Please see additional Important Safety Information on back.

Text **TRACK**
to **50334***

to start logging headaches and migraines each day—and receive a free customized report each month to bring to your doctor!



*See Privacy & Terms: <http://bit.ly/2RvxiWr>. Message & data rates may apply. Message frequency may vary. Text HELP for help or STOP to end.

“What can I expect with BOTOX[®]?”

resources to support you throughout treatment



Visit our YouTube channel for real patient stories and follow us on social media

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX[®] with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX[®] in the past.**

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as *Myobloc*[®], *Dysport*[®], or *Xeomin*[®] in the past (tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX[®] include: dry mouth, discomfort or pain at the injection site, tiredness, headache, neck pain, eye problems: double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, dry eyes; and drooping eyebrows.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see accompanying Summary of Information about BOTOX[®].

Summary of Information about BOTOX® (onabotulinumtoxinA)

What is the most important information I should know about BOTOX®?

BOTOX® may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX®:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are pre-existing before injection. Swallowing problems may last for several months
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms including: loss of strength and all-over muscle weakness, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, loss of bladder control, trouble breathing, and trouble swallowing

There has not been a confirmed serious case of spread of toxin effect away from the injection site when BOTOX® has been used at the recommended dose to treat Chronic Migraine.

BOTOX® may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of taking BOTOX®. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

BOTOX® dosing units are not the same as, or comparable to, any other botulinum toxin product.

What is BOTOX®?

BOTOX® is prescription medicine a medical professional injects into muscles to prevent headaches in adults with chronic migraine who have 15 or more days each month with headache lasting 4 or more hours each day in people 18 years and older.

It is not known whether BOTOX® is safe or effective to prevent headaches in people with migraine who have 14 or fewer headache days each month (episodic migraine).

Who should not receive BOTOX®?

Do not receive BOTOX® if you are: allergic to any of the ingredients in BOTOX® such as botulinum toxin type A and human serum albumin; had an allergic reaction to another botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), or Xeomin® (incobotulinumtoxinA); or have a skin infection at the planned injection site.

What should I tell my doctor before treatment?

Tell your doctor about all your muscle or nerve conditions, such as amyotrophic lateral sclerosis (Lou Gehrig's disease), myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects.

Tell your doctor if you have or have had breathing problems such as asthma or emphysema; swallowing problems; bleeding issues; plan to or have had surgery; have forehead muscle weakness such as trouble raising your eyebrows; drooping eyelids; or any changes to your face.

Tell your doctor if you are pregnant, plan to become pregnant, are breastfeeding or plan to breast feed. It is not known if BOTOX® (onabotulinumtoxinA) can harm your unborn baby or if BOTOX® passes into breast milk.

What Are Common Side Effects?

The most common side effects include neck pain; headache; migraine; slight or partial facial paralysis; drooping eyebrows; eyelid drooping; bronchitis; musculoskeletal stiffness; muscular weakness; pain in 1 or more muscles, ligaments, tendons, or bones; muscle spasms; injection site pain; and high blood pressure. Other side effects have been reported including allergic reactions e.g. itching, rash, red itchy welts, wheezing, asthma symptoms, or dizziness or feeling faint.

These are not all of the possible side effects. Call your doctor for medical advice if you experience any side effects after treatment with BOTOX®.

What Should I Tell My Doctor About Medicines and Vitamins I Take?

Using BOTOX® with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received BOTOX® in the past.** Tell your doctor if you have received an injection with another botulinum toxin product in the last 4 months, such as Myobloc®, Dysport®, or Xeomin®. Be sure your doctor knows which product you received.

Tell your doctor about all prescription and over-the-counter medicines, vitamins and herbal supplements you take; recent antibiotic injections; anticholinergics; muscle relaxants; allergy or cold medicine; sleep medicine; aspirin-like products; and blood thinners. **Ask your doctor if you are not sure whether your medicine is listed above.**

To Learn More

If you would like more information, talk to your doctor and/or go to BotoxChronicMigraine.com for full Product Information.

You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Based on v2.0MG1145 Rev. 06/2019

BCM69906-v4 05/20

All trademarks are the property of their respective owners. © 2020 Allergan. All rights reserved. ® marks owned by Allergan, Inc.

Patented. See: www.allergan.com/products/patent_notices

Myobloc® is a registered trademark of Solstice Neurosciences, Inc.

Dysport® is a registered trademark of Ipsen Biopharm Limited Company.

Xeomin® is a registered trademark of Merz Pharma GmbH & Co KGaA

